

1. General Information

Id 26741-53-7

Date 12.12.2003

201-15676B

I U C L I D

Data Set

Existing Chemical	:	ID: 26741-53-7
CAS No.	:	26741-53-7
EINECS Name	:	2,4,8,10-Tetraoxa-3,9-diphosphaspiro[5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl) phenoxy]-
EC No.	:	247-952-5
Molecular Formula	:	C33H50O6P2
Status	:	
Memo	:	US HPV ULTRANOX 626 Crompton Corp
Printing date	:	12.12.2003
Revision date	:	
Date of last update	:	12.12.2003
Number of pages	:	25
Chapter (profile)	:	Chapter: 1, 2, 3, 4, 5, 6, 7, 8, 10
Reliability (profile)	:	Reliability: without reliability, 1, 2, 3, 4
Flags (profile)	:	Flags: without flag, confidential, non confidential, WGK (DE), TA-Luft (DE), Material Safety Dataset, Risk Assessment, Directive 67/548/EEC, SIDS

2. Physico-Chemical Data

Id 26741-53-7

Date 12.12.2003

2.1 MELTING POINT

Value : 173 - 180 °C
Sublimation :
Method : OECD Guide-line 102 "Melting Point/Melting Range"
Year : 2003
GLP : yes
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Product Name: U626
Lot No.: H42265
Result : No color change was observed during melting point determinations, which was consistent with U626 being chemically stable at the melting point.
Reliability : (1) valid without restriction
12.12.2003 (14)

2.2 BOILING POINT

Value : > 311 °C at 1015 hPa
Decomposition : no
Method : OECD Guide-line 103 "Boiling Point/boiling Range"
Year : 2003
GLP : yes
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: U626
Lot No. H42265
Remark : Estimation using MPBPWIN v 1.40 (US EPA., EPIWIN v 3.10, EPI Suite Software, 2000) gives a boiling point of 595°C
Result : The duplicate U626 warming curves did not show a distinct boiling plateau up to the maximum temperature of 312°C (at 101.5 kPa). This was supported by the fact that U626 was not observed to boil during the test. No U626 color change nor smoke was observed during the test.
Reliability : (1) valid without restriction
12.12.2003 (13)

2.4 VAPOUR PRESSURE

Value : .0000000000029 hPa at 25 °C
Decomposition :
Method : other (calculated): MPBPWIN v 1.40
Year : 2003
GLP :
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Test condition : Melting point = 173°C (experimental)
Boiling Point = 595 (estimated)

2. Physico-Chemical Data

Id 26741-53-7

Date 12.12.2003

Reliability : (2) valid with restrictions
12.12.2003

(16)

2.5 PARTITION COEFFICIENT

Partition coefficient : octanol-water
Log pow : 10.9 at °C
pH value :
Method : other (calculated): KOWWIN v 1.66
Year : 2003
GLP :
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7

Reliability : (2) valid with restrictions
12.12.2003

(16)

2.6.1 SOLUBILITY IN DIFFERENT MEDIA

Solubility in : Water
Value : at °C
pH value :
concentration : at °C
Temperature effects :
Examine different pol. :
pKa : at 25 °C
Description :
Stable :
Deg. product :
Method : other: calculated using WSKOW v 1.40
Year : 2003
GLP :
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7

Remark : An OECD 105 study is currently underway to fill this endpoint.
Result : Water solubility = 5.67E-8 mg/L
Test condition : Melting point = 173°C (experimental)
Log Kow = 10.9 (estimated)

Reliability : (2) valid with restrictions
12.12.2003

(16)

5. Toxicity

Id 26741-53-7
Date 12.12.2003

3.1.1 PHOTODEGRADATION

Type : air
Light source :
Light spectrum : Nm
Relative intensity : based on intensity of sunlight
INDIRECT PHOTOLYSIS
Sensitizer : OH
Conc. of sensitizer : 1500000 molecule/cm³
Rate constant : .000000000000011 cm³/(molecule*sec)
Degradation : % after
Deg. product :
Method : other (calculated): estimation using AOPWIN v1.90
Year : 2003
GLP :
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Result : T1/2 = 1.166 hours
Reliability : (2) valid with restrictions
08.12.2003

(16)

3.1.2 STABILITY IN WATER

Type : Abiotic
t1/2 pH4 : at °C
t1/2 pH7 : at °C
t1/2 pH9 : at °C
Deg. product : No
Method : OECD 111
Year : 2004
GLP :
Test substance : 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Purity: >96%
Source: General Eletrical
Method : The test substance was added to pH 4, 7 and 9 buffered water and incubated at 25°C. Samples were analyzed after Day 1, 2, 3, 4, 6, 8 and 10 and analyzed by HPLC.
Result : The test substance was practically insoluble in water and difficult to determine the hydrolytic half-life. However, it is expected to be hydrolytically unstable similar to other organophosphite class of chemicals at all pHs'.
Fugacity Model Half-Life: 3.6 e003 hr
Reliability : (1) valid without restrictions
10.21.2004

(14)

5. Toxicity

Id 26741-53-7

Date 12.12.2003

3.3.1 TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS

Type : fugacity model level III
Media :
Air : % (Fugacity Model Level I)
Water : % (Fugacity Model Level I)
Soil : % (Fugacity Model Level I)
Biota : % (Fugacity Model Level II/III)
Soil : % (Fugacity Model Level II/III)
Method : other: EPIWIN Level III Fugacity Model
Year : 2003
Test Substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Test condition : Henry's Law Constant: 3.24E-9 atm-m3/mol (Henrywin)
Vapor pressure: 2.9E-12 mmHg (Mppbwin)
Melting Point: 173°C (experimental)
Log Kow: 10.9 (Kowwin)
Soil Koc: 3.26E10 (calc by model)

	Mass Amount (percent)	Half-life (hr)	Emissions (kg/hr)
Air	0.0175	2.33	1000
Water	1.25	3600	1000
Soil	33.1	3600	1000
Sediment	65.6	14400	0

	Fugacity (atm)	Reaction (kg/hr)	Advection (kg/hr)	Reaction (percent)	Advection (percent)
Air	1.12E-16	881	29.6	29.4	0.988
Water	1.07E-19	40.8	212	1.36	7.06
Soil	2.14E-21	1080	0	36	0
Sediment	1.9E-19	535	222	17.8	7.41

Persistence time: 5640 hr

Reaction time: 6680 hr

Advection time: 36500 hr

Percent reacted: 84.5

Percent advected: 15.5

Half-lives (hr), (based upon Biowin (ultimate) and Aopwin):

Air: 2.33

Water: 3600

Soil: 3600

Sediment: 14400

Biowin estimate: 0.802

Advection times (hr):

Air: 100

Water: 1000

Sediment: 5E+4

Reliability : (2) valid with restrictions

08.12.2003

5. Toxicity

Id 26741-53-7

Date 12.12.2003

3.5 BIODEGRADATION

Type : aerobic
Inoculum :
Deg. product :
Method : other: Estimation using BLOWIN v4.00
Year : 2003
GLP :
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7

Result : MITI Linear Biodegradation Probability: -0.6156
MITI Non-linear Biodegradation Probability: 0.000

Reliability : The substance is predicted to be not readily biodegradable
12.12.2003 : (2) valid with restrictions (16)

4.1 ACUTE/PROLONGED TOXICITY TO FISH

Type : other: estimation
Species :
Exposure period : 96 hour(s)
Unit : mg/l
LC50 : 1.93E-6
Method : other: calculated using ECOSAR v 0.99g
Year : 2003
GLP :
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7

Reliability : (2) valid with restrictions
12.12.2003 (16)

4.2 ACUTE TOXICITY TO AQUATIC INVERTEBRATES

Type : other: estimation
Species : Daphnia sp. (Crustacea)
Exposure period : 48 hour(s)
Unit : mg/l
EC50 : 3.82E-6
Method : other: calculated using ECOSAR v 0.99g
Year : 2003
GLP :
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7

Reliability : (2) valid with restrictions
12.12.2003 (16)

5. Toxicity

Id 26741-53-7
Date 12.12.2003

4.3 TOXICITY TO AQUATIC PLANTS E.G. ALGAE

Species :
Endpoint :
Exposure period : 96 hour(s)
Unit : mg/l
EC50 : 3.99E-6
Method : other: calculated using ECOSAR v 0.99g
Year : 2003
GLP :
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Reliability : (2) valid with restrictions
12.12.2003 (16)

5.1.1 ACUTE ORAL TOXICITY

Type : LD50
Value : 5580 mg/kg bw
Species : rat
Strain : Sprague-Dawley
Sex : male/female
Number of animals : 100
Vehicle :
Doses : 1300 - 15300 mg/kg
Method : other:IBTL method
Year : 1975
GLP : No
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Product Name: Ultranox 626 (Weston 6140)
Reliability : (4) not assignable
12.12.2003 (4)

Type : LD50
Value : > 6080 mg/kg bw
Species : hen
Strain : Leghorn
Sex : female
Number of animals : 24
Vehicle : other: corn oil
Doses : 800, 1200, 1800, 2700, 4050, 6080 mg/kg bw
Method : other: Biodynamics Inc method
Year : 1980
GLP : No

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Test substance	: Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]- CAS No.: 26741-53-7 Trade name: Weston 1452 Purity: 99.5% Lot No.: PP-357
Method	: No of animals/sex/dose: 4 hens/dose Vehicle: Corn oil Route of administration: Gavage
Result	: Number of deaths at each dose level: No hens died during the study Clinical signs: Occasional hens showed green or yellow discoloration of the feces on the day of dosing. This reflects the use of corn oil vehicle and is not considered to indicate a response to treatment. There were no signs of toxicity and the hens remained normal in respect of appearance, mood, locomotor function and body weight throughout the observation period. Necropsy findings: Occasional findings of liver discoloration . This is a common pathological entity in hens of this age and is not considered to reflect any delayed or residual response to treatment.
Test condition	: Age: 12 months Weight: 1.3 - 1.9 kg Volume administered: 10 mL/kg Post dose observation period: 14 days
Reliability 08.12.2003	: (2) valid with restrictions

(1)

5.1.2 ACUTE INHALATION TOXICITY

Type	: LC50
Value	: > 2000 mg/m ³
Species	: rat
Strain	: Sprague-Dawley
Sex	: male/female
Number of animals	: 10
Vehicle	:
Doses	: 2 mg/L
Exposure time	: 1 hour(s)
Test Substance	: Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]- CAS No.: 26741-53-7 Trade name: weston 61440
Result	: None of the animals died and all the animals appeared normal during the 14 day observation period. The gross pathological autopsy of all 10 animals showed normal appearance of the organs of the thorax and the abdomen.
Reliability 12.12.2003	: (4) not assignable

(11)

5.1.3 ACUTE DERMAL TOXICITY

Type	: LD50
Value	: > 2000 mg/kg bw

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Species : rabbit
Strain : New Zealand white
Sex : male/female
Number of animals : 10
Vehicle :
Doses : 2000 mg/kg bw
Method : OECD Guide-line 402 "Acute Dermal Toxicity"
Year : 1994
GLP : yes
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: Ultrinox 626
Lot No.: HCC033

Result : Mortality: There were no deaths during the study

Clinical observations: 5 rabbits had abnormal defecation (soft stool, mucoid feces) on day 0 or 1. One of these animals also had wet, and subsequently, dry brown urogenital matting. These findings were considered to be a result of the bandage/restraint procedures used and not related to the test material. An additional spontaneous occurrence of soft stool was noted on day 9 for one animal. There were no other findings.

Dermal observations: The test material induced generally very slight to slight erythema on all rabbits and very slight to slight edema on 8 rabbits. There was a single occurrence of moderate erythema at the day 1 observation. Desquamation was present on five sites by day 7. There were no other dermal findings. One site had very slight erythema at study termination (day 14).

Body weights: There were no remarkable changes or differences noted in body weights during the study.

Necropsy: Accessory splenic tissue, a common congenital abnormality in this strain of rabbit, was noted for four animals at the terminal necropsy. There were no other gross necropsy findings for all examined tissues.

Test condition : Age: Approximately 11 weeks old
Weight: 2014 - 2224 g at study initiation
Post dose observation period: 14 days
Reliability : (1) valid without restriction
Guideline study conducted to GLP

28.11.2003

(10)

5.2.1 SKIN IRRITATION

Species : rabbit
Concentration : .5 g
Exposure : Occlusive
Exposure time : 24 hour(s)
Number of animals : 6
Vehicle : physiol. saline
PDII :
Result : corrosive

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Classification :
Method : other: 16CFR 1500.42
Year : 1981
GLP : no
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: Weston XP-1532
Lot No.: 1198
Result : Necrotic skin occurred in all animals at 24 and 72 hours. The responses for abraded and intact skin were the same.
Reliability : (2) valid with restrictions
12.12.2003 (3)

5.3 SENSITIZATION

Type : other: Footpad method
Species : guinea pig
Concentration : 1st. Induction 1 % other: injected into footpad
2nd. Challenge 10 % open epicutaneous
3rd.
Number of animals : 20
Vehicle : other: Induction: Freund's adjuvant, Challenge: acetone/dioxane/guinea pig fat 7:2:1
Result : not sensitizing
Classification : not sensitizing
Method : OECD Guide-line 406 "Skin Sensitization"
Year : 1992
GLP : yes
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: Ultrinox 626
Lot No.: U626-1
Result : Primary irritation screen: No signs of erythema or edema were evident for the animals that were administered 1, 3 or 10% of the test material.

Sensitization study: No signs of erythema or edema were noted at the 24 or 48 hour for animals previously induced with either Freund's adjuvant (control animals) or with 1% of the test material in Freund's adjuvant).

No toxic effects or systemic clinical signs were noted during the study.
Test condition : All animals gained weight normally during the study.
Strain: Crl:(HA)BR VAF/PlusT

Primary irritation screen:

No. of animals: 5/dose
Doses: 1.0%, 3.0%, 10.0%
Sex: Not determined
Body weight range: 571 - 741 g
Age: Not determined

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Reliability : Sensitization study:
No. of animals: 20; 10 control, 10 test
Sex: Female
Body weight range: 361 - 442 g
Age at study initiation: 6 - 7 weeks
(1) valid without restriction
Guideline study conducted to GLP

28.11.2003

(15)

5.4 REPEATED DOSE TOXICITY

Type :
Species : Rat
Sex : male/female
Strain : other: Charles River CD
Route of admin. : oral feed
Exposure period : 90 days
Frequency of treatm. : Daily
Post exposure period : None
Doses : 0, 100 ppm; 300 ppm ; 1000 ppm
Control group : yes, concurrent vehicle
NOAEL : 300 ppm (Males: 22 mg/kg/bw; Females 26 mg/kg bw)
Method : other: IRDC method

The test substance was fed in the diet to three groups of 20 male and 20 female rats. The rats were observed daily for signs of overt toxicity and mortality. Detailed observations including incidence, size and location of palpable masses and individual body weights were recorded weekly. Individual food consumptions were recorded daily. All rats received the ophthalmic examinations during the pretest period and at 3 months of study. Clinical laboratory test were performed for 10 rats/sex/group at 1 and 3 months of study. At the termination of the study period, selected organ weights from 10 male/10 female rats were recorded and tissues (i.e., adrenals, aorta, eye, stomach, liver, spleen, pancreas, urinary bladder, sternum, prostate/uterus, testes/ovaries, brain, heart, lung, pituitary, mammary gland, thymus, kidneys, etc) were examined.

Year : 1979
GLP : No
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: Weston XR-1532
Lot No.: 225-35

Result : Impurities: 1% tris-isopropanolamine
Body weight: The group mean body weights of both the low and mid dose male and female rats were greater than control values, and both groups of high dose males and females had group mean body weights lower than controls.

Food/water consumption: Low and mid dose male rats consumed slightly more food than did controls, while high dose males ate slightly less than control. All of the groups of females consumed approximately equivalent amounts of food.

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Test condition

Clinical signs: No signs of overt toxicity were observed among the treated rats. Incidental signs observed for a few control and/or treated rats included hair loss, missing or malaligned upper incisor, excessive lacrimation, red material around the eyes, leaning to left, red or swollen eyes, rales, corneal opacity, swollen ventral neck, swollen conjunctiva and dilated pupil.

Ophthalmologic findings: There were no compound related effects noted during the 3-month ophthalmic examinations.

Hematology: No compound related effects on the results of the hematologic tests were observed.

Clinical biochemistry: No compound related effects on the results of the biochemical tests were observed.

Mortality: No compound related effects on survival rates were noted (one female rat in the 1000 ppm group died during the study).

Gross pathology: No compound related gross lesions were observed in any of the rats from the treated groups. No tumors were noted upon gross examination.

Organ weight changes: Statistically significant variations ($p < 0.01$) in absolute weights of kidneys of male rats at 300 ppm and hearts of male rats at 100 and 300 ppm were not considered compound related.

Histopathology: Microscopic lesions considered probably compound related were seen in livers and spleens of the female rats from the 1000 ppm group. This consisted of very slight to slight extramedullary hematopoiesis in these organs. This lesion was not present in rats from the control and the 300 ppm groups but was seen in one rat from the 100 ppm group. Other microscopic lesions in livers and those seen in other organs in the control and 1000 ppm groups were considered spontaneous in nature and unrelated to the administration of the compound.

: **Statistical methods:** All statistical analyses compared the treatment groups with the control group by sex. Body weights, food consumption, absolute and relative organ weights and hematology, biochemistry and urinalysis parameters were compared by analysis of variance (one-way), Bartlett's test for homogeneity of variances and the appropriate t-test (for equal or unequal variances) as described by Steel and Torrie using Dunnett's multiple comparison tables to judge significance of differences.

Test subjects:

Weight at study initiation: 80-107g (male), 77-98g (female)

No. of animals/sex/dose: 20 male, 20 female

Study Design:

Vehicle: Feed (Rodent Laboratory Chow)

Clinical observations performed and frequency: The rats were observed twice daily for signs of overt toxicity and mortality. Detailed observations were recorded weekly and included the size, incidence and location of all

5. Toxicity

Id 26741-53-7

Date 12.12.2003

palpable masses. Individual body weights were recorded weekly and individual food consumption was recorded daily.

Organs examined at necropsy: The following tissues from 10 males and 10 females from the control group and the 1000 ppm group were examined microscopically: adrenals, aorta, eye and optic nerve, esophagus, stomach, duodenum, jejunum, ileum, cecum, colon, rectum, liver, trachea, spleen, pancreas, urinary bladder, bone marrow (sternum), prostate/uterus, seminal vesicles, testes/ovaries, brain, heart, lung and bronchi, sciatic nerve, pituitary, thyroid and parathyroid, mesenteric lymph node, mandibular lymph node, spinal cord, salivary gland (submaxillary), skeletal muscle (thigh), skin, mammary gland, thymus, kidneys, any other tissue with gross lesions.

Bone marrow smears from all rats were made at necropsy and examined microscopically.

Additionally, livers and spleens from 10 male and 10 female rats from the 100 ppm and 300 ppm groups were examined.

Reliability : (2) valid with restrictions
Well conducted study, prior to GLP

05.12.2003

(9)

Type :
Species : Dog
Sex : male/female
Strain : Beagle
Route of admin. : other: oral via gelatin capsule
Exposure period : 4 months
Frequency of treatm. : daily
Post exposure period :
Doses : 0, 4, 12, 40 mg/kg/day
Control group : yes
NOAEL : 12 mg/kg bw
Method : other: Bio/dynamics Inc method
Year : 1980
GLP : no
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: Weston XR-1532
Lot No.: 1532-003-04189-X
Purity: 100%

Result : Mortality: One high dose female was sacrificed in a moribund condition after 86 days on test. All of the remaining control and treated animals survived the duration of the study.

Physical and neurological observations: The only physical and neurological observations noted during the study which were considered related to the administration of the test substance involved the high dose female which was sacrificed in a moribund condition. On test day 51, this animal exhibited the first signs of uncoordination of the hind limbs and a poor righting reflex. These symptoms progressed and by test day 58, the animal displayed a lack of coordination of both the fore and hind limbs. The

5. Toxicity

Id 26741-53-7

Date 12.12.2003

animal became progressively recumbent as the uncoordination progressed to rigid fore and hind limb paralysis. During this time the animal was offered canned dog food in an attempt to increase its food consumption and maintain its body weight. However this was unsuccessful and the animal was sacrificed in a moribund condition on test day 86.

Ophthalmology: There were no ocular abnormalities observed after 3 or 4 months of study which could be attributed to the test substance.

Body weight and food consumption: The mean body weights of all treated males were slightly greater than control prior to initiation of dosing (4-5%) and throughout the 4 month treatment period. At 18 weeks these differences from control were 25%, 20%, and 21% in the low, mid and high dose males, respectively. Mean food consumption values were slightly lower in the low (13%), mid (12%) and high (17%) dose males when compared to the control values over the 18 week period of test substance administration. Mean body weight and food consumption values for the control and treated females were unremarkable throughout the study with the exception of one high dose female. During week 9 this animal began to exhibit decreases in food consumption and body weight. These decreases continued over the next 3 weeks and the animal was sacrificed in a moribund condition on test day 86.

Hematology and clinical chemistry: At 1 month the mean platelet counts of the treated males were significantly lower than control. However, these differences were attributed to a slightly greater mean platelet counts in the control group rather than an effect of the test substance. Other differences from control were noted in some of the hematology and clinical chemistry parameters evaluated. However, these differences were not dose related or consistent over time and therefore were not considered related to administration of the test material.

Urinalysis: The urinalysis data collected after 1, 3 and 4 months of study were unremarkable and revealed no differences between control and treated groups which could be attributed to the administration of the test substance.

Organ weights and organ/body weight ratios: Slight differences from control, some statistically significant, were observed in the mean absolute and relative organ weights of the treated animals. These variations were attributed to differences in terminal body weights and normal biological variation between animals. There were no differences observed in absolute or relative organ weights which were attributed to the administration of the test substance.

Pathology: A subacute, eosinophilic pneumonia was observed microscopically in four of eight high dose animals, one of eight mid dose animals and two of eight low dose animals. This condition was not observed in any of the control animals. The etiology of the pulmonary lesions could not be determined. Seven of the eight high dose animals displayed degenerative myelin lesions. These lesions were confined to the high dose group and were considered related to the administration of the test material. One animal (the female which died) displayed clinical manifestations of several abnormalities of the axonal fibers and myelin.

Test condition

: Statistical methods: Hematology and clinical chemistry: Test substance groups were compared to control by the F-test and Student's T-test. When

5. Toxicity

Id 26741-53-7

Date 12.12.2003

variances differed significantly (F-test), Student's t-test was appropriately modified using Cochrane's approximation (t'). Snedecor, G.W. & Cochran, W.G. (1967) Statistical Methods, 6th ed. Iowa State University Press, Ames, pp 104-106, 114-119. Body weight, Food consumption, Organ weights and organ/body weight ratios: Treated groups were compared to control. Dunnett, C.W. (1964) J. Am. Stat. Assn. 50, 1096-1121 and Biometrics 20, 482 (1964).

Test subjects:

Weight at study initiation: 80-107g (male), 77-98g (female)

No. of animals/sex/dose: 4/sex/dose

Study Design:

Vehicle: None

Clinical observations performed and frequency: Twice daily for mortality and gross signs of toxicological or pharmacologic effects.

Detailed physical examination: Pretest and weekly thereafter

Neurologic examination: Pretest and monthly thereafter

Ophthalmoscopic examination: Pretest, 3 and 4 months

Body weight: Pretest, weekly during treatment and terminally

Food consumption: Pretest and weekly thereafter

Laboratory studies: Pretest and monthly thereafter. Parameters evaluated - Hematology: hemoglobin, hematocrit, erythrocytes, platelets, clotting time, prothrombin time, total and differential leukocytes, erythrocyte morphology. Clinical chemistry: serum glutamic oxaloacetic transaminase, serum glutamic pyruvic transaminase, alkaline phosphatase, blood urea nitrogen, fasting glucose, total protein, albumin, globulin, A/G ratio, cholesterol, triglycerides, sodium, potassium, chloride, calcium, creatinine, lactic acid dehydrogenase, total bilirubin, direct bilirubin. Urinalysis: gross appearance, specific gravity, pH, protein, glucose, ketones, bilirubin, occult blood, microscopic analysis

Organs examined at necropsy: Organs weighed and organ/body weight ratios calculated: adrenals, brain, ovaries, testes (without epididymides), heart, kidneys, liver (with gall bladder attached), pituitary, spleen, thyroid (with parathyroid). Tissues examined histopathologically: adrenal, aorta (thoracic, arch, lumbar), bone and bone marrow (sternum), brain, epididymus, esophagus, eye, heart, intestine, cecum, colon, rectum, kidney, liver with gall bladder, lungs with mainstem bronchi, lymph nodes (cervical, peribronchial, mesenteric), nerve (left sciatic), oral mucous membrane (including tongue, buccal, maxillary gingiva, pharynx and nasopharynx), ovary, pancreas, pituitary, prostate, salivary gland (submaxillary), skeletal muscle (biceps femoris), skin with mammary gland (left inguinal), spinal cord (cervical, thoracic, lumbar), spleen, stomach (cardia, pylorus, fundus), testis, thymus, thyroid with parathyroid, trachea,

5. Toxicity

Id 26741-53-7

Date 12.12.2003

		urinary bladder, uterus (corpus, cervix, fallopian tubes), vagina, gross lesions, tissue masses or suspect tumors and regional lymph nodes.	
Reliability	:	(2) valid with restrictions	
05.12.2003		Well conducted study, prior to GLP	(2)
Type	:		
Species	:	Rat	
Sex	:	Male/female	
Strain	:	Wistar	
Route of admin.	:	oral feed	
Exposure period	:	24 months	
Frequency of treatm.	:	Daily	
Post exposure period	:	None	
Doses	:	100, 500 ppm	
Control group	:	yes, concurrent no treatment	
NOAEL	:	500 ppm	
Method	:	other: NIHMR method	
Year	:	1983	
GLP	:	No	
Test substance	:	Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]- CAS No.: 26741-53-7 Trade name: Weston XP1452 Purity: 99% Impurities: Tri-isopropanolamine (1%)	
Result	:	Body weight: No difference was observed between the average weights of the control group and treated animals. Food/water consumption: No difference was observed between the treated groups and the control group animals. Clinical signs: No difference between the treated animals and the control group. Toward the 18th month, the animals began to show signs of ageing, i.e. some of them had a yellow coat, dyspnoea, pulmonary insufficiency and dark rings around the eyes, but these symptoms were relatively rare and no more frequent among the treated animals than among the control animals. Hematologic findings: The haemoglobin level, the haematocrit, the number of erythrocytes and the level or relative number of leucocytes were the same in all the groups. Clinical biochemistry: No difference in glucose levels, ureic nitrogen levels, the GOT, the GPT, the PA, total proteins and the albumin levels were seen. Renal function: The urine density showed no significant variation and neither did the proteinurea, glycosuria and urinary residue examinations. Mortality:	
		Dose Sex Total Months to death (mg/kg) Deaths (No. of animals)	

5. Toxicity

Id 26741-53-7

Date 12.12.2003

0	Male	11	13(1), 19(4), 20(1), 21(3), 22(2)
	Female	5	20(1), 21(2), 22(2)
100	Male	9	12(1), 13(1), 15(1), 16(1), 17(1), 20(1), 21(2), 22(1)
	Female	5	19(1), 21(3), 22(1)
500	Male	15	13(1), 14(2), 15(1), 16(1), 17(3), 18(1), 20(1), 21(1), 22(4)
	Female	6	14(2), 15(1), 21(1), 23(1)

Mortality was higher among the males than among the females. A dose of 100 ppm did not give rise to higher mortality among the male or female rats compared to the control group. At 500 ppm, mortality is no higher among the female rats, but higher mortality was observed among the male rats. The authors concluded that this difference was not significant.

Gross pathology: an examination of the main organs did not reveal any significant difference between the treated rats and the control group. The animals often showed clear organic pathological signs of aged animals which could in no case be attributed to the treatment since they appeared with equal frequency in the control group.

Organ weight changes: No change in the weight of the main organs under the effect of the treatment was seen.

Histopathology: No greater frequency of anomalies in the organs of the treated rats compared with the control group.

Test condition

: Test subjects

Age at study initiation: no data

No. of animals/sex/dose: 30 males/females per dose

Study Design

Vehicle: Feed

Clinical observations: Each animal was weighed every week over the first four weeks, twice a week during weeks 6-12 and then every other week after week 12. Food and water intake was measured in weeks 1+3, 3+4, 11+12, 23+24, 38+39, 52+53, 75+76, 86+87 and 97+98.

Haematological examinations were conducted during weeks 12, 24, 52, 88 and 104. The blood was used:

- a) to conduct measurement of haemoglobin content, haematocrits, number of erythrocytes and leucocytes and leucocytic formula;
- b) to determine blood glucose and ureic nitrogen levels;
- c) to analyze for the following enzymatic activities: glutamic-pyruvate transaminase, alkaline phosphatase and proteins and albumin serum.

Urine was collected in weeks 12, 24, 53, 98-104 and tested for appearance, pH, glucose, proteins, density, blood, lymphocytes and erythrocytes, epithelial cells, urate and phosphate crystals and evaluation of the bacterial population.

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Organs examined at necropsy: The main organs were subjected to a macroscopic examination and some were weighed (liver, kidneys, spleen, brain, heart, supra-renal bodies and hypophysis). Tissue fragments were taken: heart, kidney, liver, pancreas, spleen, brain, ovary or testicle, thyroid, suprarenal bodies, thymus, lung, trachea, salivary glands, stomach, duodenum, colon, caecum and uterus.

Reliability : (2) valid with restrictions
10.12.2003 (5)

i.p GENETIC TOXICITY 'IN VITRO'

Type : Ames test
System of testing : Salmonella typhimurium strains TA97, TA 98, TA100, TA102
Escherichia coli Strain WP2 (PKM101)
Test concentration : 50 – 2000 µg/Plate
Cycotoxic concentr. :
Metabolic activation : with and without
Result : negative
Method : other: Hachiya (1994)
Year : 1994
GLP : no data
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7

Test condition : Solvent: Acetone and Tween 80
Reliability : (4) not assignable
Secondary literature data
08.12.2003 (8)

Type : Chromosomal aberration test
System of testing : Chinese hamster ovary cells
Test concentration : 31.3-2000 µg/mL (-S9, 4 hour exposure), 31.3 – 3000 µg/mL (+S9, 4 hour exposure; -S9, 20 hour exposure)
Cycotoxic concentr. : 500 µg/mL (+S9), 2000 µg/mL (-S9)
Metabolic activation : with and without
Result : positive
Method : OECD Guide-line 473
Year : 2003
GLP : yes
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: Ultrinox 626
Lot No.: H42265

Method : METHOD

Metabolic activation: Arochlor 1254-induced rat liver S(9)

Concentrations tested:

Experiment 1:

31.3, 62.5, 125, 250, 500, 1000 µg/mL (-S9, 4 hour exposure)

31.3, 62.5, 125, 250, 500, 1000, 1500, 3000 µg/mL (-S9, 20 hour

5. Toxicity

Id 26741-53-7

Date 12.12.2003

exposure)
31.3, 62.5, 125, 250, 500, 1000, 1500, 3000 (+S9, 4 hour exposure)

Result

Statistical Methods: Statistical analysis of the percent aberrant cells was performed using Fisher's exact test. Fisher's exact test was used to compare pairwise the percent aberrant cells of each treatment group with that of the solvent control. In the event of a positive Fisher's exact test at any test article dose level, the Cochran-Armitage test was used to measure dose-responsiveness.

: +S9, 4h exposure: The percentage of cells with structural aberrations in the test article treated group was statistically increased above that of the solvent control at 250 µg/mL ($p=0.05$, Fisher's exact test). The Cochran-Armitage test was also positive for a dose response ($p<0.05$). However, the percentage of cells with structural aberrations in the test article treated group (6.0%) was within the historical solvent control range of 0.0 – 6.5%. Therefore it is not considered to be biologically significant. The percentage of cells with numerical aberrations in the test article treated groups was not significantly increased above that of the solvent control at any dose level ($p>0.05$, Fisher's exact test).

–S9, 20h exposure: The percentage of cells with structural or numerical aberrations in the test article treated groups was not significantly increased above that of the solvent control at any dose level ($p>0.05$, Fisher's exact test).

–S9, 4h exposure: Due to the lack of dose levels with $\approx 5\%$ toxicity relative to solvent control in this group, the experiment was repeated at higher dose levels.

–S9, 4h exposure (repeat): The percentage of cells with structural aberrations in the test article treated groups was significantly increased above that of the solvent control at 2000 µg/mL ($p=0.05$, Fisher's exact test). The Cochran-Armitage test was also positive for a dose related response ($p<0.05$). The percentage of cells with numerical aberrations in the test article-treated groups was not significantly increased above that of the solvent control at any dose level ($p=0.05$, Fisher's exact test).

Test condition

: Test Design

Number of replicates: Duplicate tests

Positive and negative controls and treatment: Mitomycin C was used as the positive control in the non-activated study at final concentrations of 0.1 and 0.2 µg/mL. Cyclophosphamide was used as the positive control in the S9 activated study at final concentrations of 10 and 20 µg/mL. The solvent for the test article (DMSO) was used as the solvent control at the same concentration as that found in the test article-treated groups.

Solvent: dimethylsulfoxide

Number of metaphases analyzed: Whenever possible, a minimum of 200 metaphase spreads (100 per duplicate flask) were examined and scored for chromatid-type and chromosome-type aberrations. The number of metaphase spreads that were examined and scored per duplicate flask was reduced when the percentage of aberrant cells reached a statistically significant level before 100 breaks were scored.

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Description of follow up repeat study: Due to lack of dose levels with 50% toxicity relative to the solvent control in the non-activated 4 hour exposure group, the chromosome aberration assay was repeated in that group at dose levels of 250, 500, 1000, 1250, 1500 and 2000 µg/mL.

Criteria for evaluating results: The test article was considered to induce a positive response when the percentage of cells with aberrations increased in a dose-responsive manner with one or more concentrations being statistically significant ($p=0.05$). However, values that were statistically significant but did not exceed the range of historic solvent controls was judged as not biologically significant. Test articles not demonstrating a statistically significant increase in aberrations were concluded to be negative.

Conclusion : Under the conditions of the assay described in this study, the test material was concluded to be weakly positive for the induction of structural and negative for the induction of numerical chromosome aberrations in CHO cells in the absence of metabolic activation. The test material was concluded to be negative for the induction of structural and numerical chromosome aberrations in CHO cells in the presence of metabolic activation.

Reliability : (1) valid without restriction
25.11.2003

(7)

i.p GENETIC TOXICITY 'IN VIVO'

Type : Micronucleus assay
Species : mouse
Sex : male/female
Strain : ICR
Route of admin. : i.p.
Exposure period : 24, 48 hours
Doses : 500, 1000, 2000 mg/kg
Result : negative
Method : OECD Guide-line 474 „Genetic Toxicology: Micronucleus Test“
Year : 2003
GLP : yes
Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: Ultrinox 626
Lot No.: H42265

Result : Effect on mitotic index or PCE/NCE ratio by dose level by sex: See table below

Genotoxic effects: Negative

Mortality at each dose level by sex:

Pilot toxicity study: No mortality occurred at any doses during the course of the study.

Main study: No mortality occurred at any dose during the course of the

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Test condition

micronucleus study.

Clinical signs:

Pilot toxicity study: All animals appeared normal except male mice at 2000 mg/kg, which exhibited piloerection.

Main study: Piloerection was seen in all male and female mice at 1000 and 2000 mg/kg. All other mice appeared normal during the study.

Bodyweight changes:

Pilot toxicity study: Change in group mean bodyweights ranged from -0.3% (1.0 mg/kg) to 4.5% (1000 mg/kg) after 3 days.

Mutant/aberration/mPCE/polyploidy frequency, as appropriate: See table below.

Food/water consumption: no data

: Age at study initiation: 6 – 8 weeks old at the initiation of each phase of the study.

No. of animals per dose:

Pilot toxicity study: 2 male mice dosed at 1, 10, 100 or 1000 mg/kg b.w.; 5 male and 5 female mice dosed at 2000 mg/kg.

Main study: Groups of 5 male/5 female mice dosed at 0, 500, 1000, 2000 mg/kg (euthanized at 24 h); Groups of 5 male/5 female dosed at 0, 2000 mg/kg (euthanized at 48 h).

Route: i.p.

Vehicle: Corn oil.

Controls: Vehicle (Corn oil), cyclophosphamide monohydrate (positive).

Clinical observations performed: Clinical signs, mortality, bodyweight

Organs examined at necropsy: none

Criteria for evaluating results: The incidence of micronucleated polychromatic erythrocytes per 2000 polychromatic erythrocytes was determined for each mouse and treatment group. Statistical significance was determined using the Kastenbaum-Bowman tables which are based on the binomial distribution. In order to quantify the proliferation state of the bone marrow as an indicator of bone marrow toxicity, the proportion of polychromatic erythrocytes to total erythrocytes was determined for each animal and treatment group. The test article was considered to induce a positive response if a dose-responsive increase in micronucleated polychromatic erythrocytes was observed and one or more doses were statistically elevated relative to the vehicle control ($p \leq 0.05$, Kastenbaum-Bowman Tables) at any sampling time. However, values that were statistically significant but did not exceed the range of historical negative or vehicle controls were judged as not biologically significant. The test article was judged negative if no statistically significant increase in micronucleated polychromatic erythrocytes above the concurrent vehicle control values and

5. Toxicity

Id 26741-53-7

Date 12.12.2003

no evidence of dose responses were observed at any sampling time.

Criteria for selection of M.T.D.: based on preliminary toxicity study.

Table: Summary of Bone Marrow Micronucleus analysis

Treatment (20mL/kg)	Sex	Time (hr)	No. of mice	PCE/Total Erythrocytes (mean \pm SD)	Change from Control (%)	Micronucleated Polychromatic Erythrocytes	
						Number per 1000 PCEs (mean \pm SD)	Number per PCEs Scored ¹
Corn oil	M	24	5	0.456 \pm 0.07	-	0.6 \pm 0.22	6/ 10000
	F	24	5	0.526 \pm 0.09	-	0.5 \pm 0.35	5/ 10000
Test article							
500 mg/kg	M	24	5	0.477 \pm 0.06	5	0.6 \pm 0.22	6/ 10000
	F	24	5	0.462 \pm 0.04	-12	0.6 \pm 0.42	6/ 10000
1000 mg/kg	M	24	5	0.480 \pm 0.06	5	0.7 \pm 0.27	7/ 10000
	F	24	5	0.543 \pm 0.05	3	0.8 \pm 0.27	8/ 10000
2000 mg/kg	M	24	5	0.493 \pm 0.09	8	0.5 \pm 0.00	5/ 10000
	F	24	5	0.469 \pm 0.04	-11	0.3 \pm 0.27	3/ 10000
CP ² 50 mg/kg	M	24	5	0.335 \pm 0.03	-27	22.2 \pm 2.20	*222/ 10000
	F	24	5	0.325 \pm 0.01	-38	20.4 \pm 2.43	*204/ 10000
Corn oil	M	48	5	0.502 \pm 0.06	-	0.3 \pm 0.27	3/ 10000
	F	48	5	0.483 \pm 0.05	-	0.6 \pm 0.22	6/ 10000
Test article							
2000 mg/kg	M	48	5	0.447 \pm 0.05	-11	0.6 \pm 0.22	6/ 10000
	F	48	5	0.495 \pm 0.05	2	0.5 \pm 0.35	5/ 10000

¹*statistically significant, $p \leq 0.05$ (Kastenbaum-Bowman Tables).

² cyclophosphamide monohydrate

Reliability : (1) valid without restriction
26.11.2003

(6)

5.8.1 TOXICITY TO FERTILITY

5.8.2 DEVELOPMENTAL TOXICITY/TERATOGENICITY

Species : rabbit
Sex : female
Strain :
Route of admin. : oral unspecified
Exposure period : Days 6 to 18 of gestation
Frequency of treatm. : Daily
Duration of test : Fetuses removed on day 29 of gestation
Doses : 0, 20, 50, 200 mg/kg
Control group : Yes
Method : The test material was administered to 4 groups of 15 rabbits at dose levels of either 0, 20, 50 or 200 mg/kg from days 6 to 18 of gestation. The fetuses were removed by caesarean section on day 29 of gestation and examined for malformations.
Year :
GLP : Well conducted study, prior to GLP

5. Toxicity

Id 26741-53-7

Date 12.12.2003

Test substance : Chemical name: 2,4,8,10-tetraoxa-3,9-diphosphaspiro [5.5]undecane, 3,9-bis[2,4-bis(1,1-dimethylethyl)phenoxy]-
CAS No.: 26741-53-7
Trade name: XP1452

Result : No maternal effects (i.e. body weights, clinical and pathological observations) were noted in any dose group.

3/15 rabbits miscarried in the high dose group (200 mg/kg) however, this finding was considered only bordering significance.

The number of implantations and the number and weight of the fetuses were not significantly different from the control values.

There was no difference in the distribution between male and female fetuses and there was not a significant number of malformations observed (1 fetus with aplasia of the head at 50 mg/kg and 1 fetus with internal hydrocephalus at 200 mg/kg).

No mortality was observed between the treated animals and the control group.

In four series of tests, the biological and biochemical parameters proved to be normal.

The histopathological results did not reveal any difference between the treated and the control animals.

Conclusion : Ultranox 626 is not consider the substance to be a teratogenic agent.
Reliability : (2) valid with restrictions

12.12.2003

(12)

9. References

Id 26741-53-7

Date 12.12.2003

- (1) Ben-dyke, R (1980), Biodynamics, Acute toxicity study in the chicken on Weston 1532, Project No. 6234-80
- (2) Bio/dynamics Inc. (1980), A four month oral toxicity study with Weston XR 1532 in dogs, Project No. 79-2411
- (3) Bio/dynamics Inc. (1981) Primary Dermal Irritation Study in Rabbits, Project No. 6772-81
- (4) Brett, B (1975) Industrial Bio-Test Laboratories, Acute oral toxicity study with CDP-1106 in albino rats, Report No. 601-07166; in, unpublished Toxicity Summary of Ultranox 626, prepared by GE Specialty Chemicals, 1997
- (5) Derache, R (1983) National Institute of Health and Medical Research, Long term study (2 years) on rats of the effect of the possible toxicity of a product called XP 1452, Report No. RD 3-83
- (6) Gudi, R., & Krsmanovic, L. (2003) Bioreliance, Mammalian erythrocyte micronucleus test, Study No. AA77XB.123.BTL
- (7) Gudi, R., & Rao, M., (2003) Bioreliance, In vitro mammalian chromosome aberration test, Study No. AA77XB.331.BTL
- (8) Hachiya, N & Takizawa, Y (1994) Mutagenicity of plastic additives, Hen'Igensei Shiken 3(3), 147-154; reported in the Chemical Carcinogenesis Research Information System database
- (9) International Research and Development Corporation (1979), 90 day oral toxicity study in rats, Report No. 433-001
- (10) Kiplinger, GR (1994), Acute dermal toxicity study of Ultranox U626 in albino rabbits, Project No. WIL-202007
- (11) Levi, RS (1978) Kemron Environmental Services, Report No. 80600; in, unpublished Toxicity Summary of Ultranox 626, prepared by GE Specialty Chemicals, 1997
- (12) Paul Sabatier University, National Institute of Health and Medical Research, Study on teratogenicity in the 'Grand Fauve De Bourgogne' rabbit after oral administration, Report No. RD 3-83; in, unpublished Toxicity Summary of Ultranox 626, prepared by GE Specialty Chemicals, 1997
- (13) Reimer, GJ (2003) BC Research Inc., Physical/Chemical Property of U626; CAS # 26741-53-7, Boiling Point (OECD 103), Study No. RAA6233 1257-BP
- (14) Reimer, GJ (2003) BC Research Inc., Physical/Chemical Property of U626; CAS # 26741-53-7, Hydrolytic Stability (OECD 111), Study No. RAA6233 1257-Hy
- (15) Shepard, KP (1992) Toxicological Sciences Laboratory (Eastman Kodak), Bis (2,4-di-t-butylphenyl) pentaerythritol diphosphite; Synonyms: Ultranox 626; PM 5753; skin sensitization study (Footpad method) in the Guinea Pig, HAEL No. 92-0046
- (16) US EPA., EPIWIN v 3.10, EPI Suite Software, 2000